

BENEFITS OF IMPLANTS

•A free-standing tooth replacement:

Unlike a bridge, implants do not involve the adjacent teeth for support. This means you would not need to shave down adjacent healthy teeth to support a bridge. Also, if something goes wrong with a bridge, at least 3 teeth would need to be treated to fix the problem, whereas with an implant, only the one tooth space is dealt with.



Bridge: Adjacent teeth are shaven down to support the bridge

•Ease of cleaning:

Unlike a bridge which can't be flossed normally, implants can be treated more like natural teeth using simply a toothbrush and floss for maintenance. Occasionally a proxy-brush or Waterpik may be useful.

• Long term restoration:

The implant fixture should remain in your bone for the rest of your lifetime, PROVIDED you clean it daily, just like the rest of your teeth!

RISKS OF TREATMENT

- Temporary Bleeding, Swelling, Bruising, Infection and/or Pain.
- Infection that is chronic leading to implant loss.
- Possible Permanent Numbness in the lower lip.


DENTAL IMPLANTS AND MISSING TEETH

Dental implants are the most advanced method we have in dentistry today to replace missing teeth. They were first used in 1978, and have made great advances over the years. As long as you have stopped growing, young and old alike can use dental implants to replace missing teeth.

Implants are metal screws made of titanium (the same material used in hip and knee replacements) which are placed into the bone to replace a missing tooth or a group of missing teeth. Titanium has been shown to make a special bond with bone called osseointegration. It takes at least 6 weeks for this process to be completed so that the implant is integrated to the bone.

After osseointegration, the crown or bridge, or even a denture, usually connected to the implant by a piece called the abutment, can be fitted to the implants by your dentist so you can begin to chew and function with them. Sometimes it takes longer than 3 months for the implant to integrate to the bone, de-

pending on the quality of bone that the implant will be placed into, or if there is a bone graft or sinus graft placed at the same time.




WHAT IF I DO NOTHING

When teeth are missing, adjacent and opposing teeth can drift into the wrong position. This can disrupt esthetics, jaw joint comfort, chewing function, and cause excessive wear or load on remaining teeth leading to their early loss.

Copyright 2019 Dr. Pamela Nicoara

Pamela A Nicoara DDS MSD PLLC

PERIODONTOLOGY IMPLANTOLOGY ORAL MEDICINE



3125 Colby Avenue, Suite H
Everett WA 98201
T: 425-374-5380 F: 425-374-5382

www.NICOARAPERIO.com
doctor@NICOARAPERIO.com

PATIENT PAGES

A SURGICAL PERIODONTAL BROCHURE FOR PATIENTS

BY DR. PAMELA NICOARA

Dental Implants for Front Teeth



Surgical Considerations

DAY OF SURGERY

- Expect about 1 hr for surgery for 1 tooth (more time needed for more teeth).
- Photos of the gums only will be taken before and after to record initial appearance and outcomes.
- Local anesthetic ('Novocaine' to numb the area).
- Sedation if desired (must be organized prior to the day of surgery) in 2 forms below. You will need a driver to bring you and take you home who can speak and read English and sign the informed consent for you. You may not drive, take a taxi or bus, or leave alone.
 1. Halcion: Anti-anxiety Tablet with sleepiness and amnesia as side effects. or
 2. General Sedation: with an anesthesiologist to 'knock you out'.
- Don't wear any make up.
- Eat a light breakfast.
- Bring headphones /iPod if you want.
- No work or exercise the rest of that day, and possibly for the day or two after especially if tissue is taken from the palate and your job is highly physical (i.e.: FedEx delivery person) or requires a lot of talking (i.e.: receptionist)

PATIENT RESPONSIBILITIES

Follow all instructions provided: take medications (antibiotics and/or pain killers) on time, use the mouth rinse, do NOT touch or eat on the areas treated, rest.

I have read this brochure and understand what may be involved in my treatment. I will have the opportunity to ask questions prior to my surgery.

Patient Signature

Date

Dental Implants

IMPORTANT ANATOMY

Implants require bone to surround the entire screw. When teeth are lost, the bone naturally shrinks away. Rebuilding that bone requires bone grafting. Sometimes the bone graft can be done at the same time the implant is placed, and sometimes so much bone is missing, that the graft has to be done first (see below).



Pre-Graft and
3 months Post-Graft

SINGLE TOOTH REPLACEMENT

Sometimes simply placing an implant is all that is needed to replace a missing tooth, and there is no bone grafting needed so the process is relatively

Single Tooth Replacement



simple. Most times, however, because tissues don't improve with time, bone and gum grafting is necessary. Sometimes, adjacent teeth need to be treated to achieve optimal esthetic results. For Dr. Nicoara's patient in the prior panel, the tooth replaced with an implant had bone and gum grafting to reduce the recession, while the tooth adjacent had gum and bone trimmed away to level the gum margins. New crowns were placed on the teeth adjacent to the implant in order to match the implant crown and improve tooth colors and shapes to satisfy the patient's esthetic ideals.

THE PROCESS

Implants can sometimes be a time consuming process, but the long-term benefits generally outweigh this disadvantage.


Typically, the final crown is not placed on the implant the same day the implant is placed (usually 3-4 months of healing is required), and a temporary tooth replacement in the form of either an *'essix appliance'* or *'flipper'* is used to help hide the fact that the tooth is missing. For highly esthetic cases in the front of the mouth, our office will fabricate a *temporary implant*



Essix



Flipper



crown made of plastic similar to the final crown. However, it has a very specific form used for shaping the tissues into their most ideal contours before your dentist makes the final crown. Of the two patients below, you can identify that the temporary crown on the right is fitting much better, and the gums look much more natural and healthy around it, than the final crown on the left. The more natural the tissue contours, the easier the implant will be to clean and maintain for the rest of your lifetime.




Final Crown

Temporary Crown

MAINTENANCE

Implants, similar to teeth, can experience bone loss, called *'Peri-Implantitis'*. Regular cleanings and your own daily hygiene can prevent such bone loss from occurring. Special surgical techniques can be used to attempt to arrest and sometimes regenerate bone around an implant, but if bone loss is severe enough, the implant cannot be saved and must be removed.



Healthy Implant



Peri-Implantitis

Visit our website: www.NICOARAperio.com